


Methocarbamol



$C_{11}H_{15}NO_5$ 241.24
1,2-Propanediol, 3-(2-methoxyphenoxy)-, 1-carbamate,
(±)-;
(±)-3-(*o*-Methoxyphenoxy)-1,2-propanediol 1-carbamate
[532-03-6].

DEFINITION

Methocarbamol contains NLT 98.5% and NMT 101.5% of methocarbamol ($C_{11}H_{15}NO_5$), calculated on the dried basis.

IDENTIFICATION

- **A. INFRARED ABSORPTION** (197K)

Change to read:

- **B.** ▲The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.▲^{USP38}

ASSAY

Change to read:

- **PROCEDURE**

▲**Buffer:** 6.8 g/L of monobasic potassium phosphate in water. Adjust with phosphoric acid or sodium hydroxide to a pH of 4.5.

Mobile phase: Methanol and *Buffer* (30:70)

System suitability solution: 1.0 mg/mL of USP Methocarbamol RS and 0.005 mg/mL of USP Guaifenesin RS in *Mobile phase*

Standard solution: 0.1 mg/mL of USP Methocarbamol RS in *Mobile phase*

Sample solution: 0.1 mg/mL of Methocarbamol in *Mobile phase*

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 274 nm

Column: 4.6-mm × 15-cm; 3- μ m packing L1

Column temperature: 30°

Flow rate: 0.8 mL/min

Injection volume: 20 μ L

Run time: 1.5 times the retention time of methocarbamol

System suitability

Samples: *System suitability solution* and *Standard solution*

[NOTE—See *Table 1* for relative retention times.]

Suitability requirements

Resolution: NLT 3.5 between methocarbamol and guaifenesin, *System suitability solution*

Tailing factor: NMT 2.0, *Standard solution*

Relative standard deviation: NMT 0.73%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of methocarbamol ($C_{11}H_{15}NO_5$) in the portion of Methocarbamol taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of methocarbamol from the *Sample solution*

r_S = peak response of methocarbamol from the *Standard solution*

C_S = concentration of USP Methocarbamol RS in the *Standard solution* (mg/mL)

C_U = concentration of Methocarbamol in the *Sample solution* (mg/mL)▲^{USP38}

Acceptance criteria: 98.5%–101.5% on the dried basis

IMPURITIES

- **RESIDUE ON IGNITION** (281): NMT 0.1%

Delete the following:

- **HEAVY METALS, Method I** (231)

Sample solution: 1.0 g in a 10-mL mixture of methanol and 1 N acetic acid (7:3), diluted with water to 25 mL

Acceptance criteria: NMT 20 ppm● (Official 1-Dec-2015)

Change to read:

- **ORGANIC IMPURITIES**

▲**Mobile phase, System suitability solution, and Chromatographic system:** Proceed as directed in the *Assay*.

Standard solution: 0.005 mg/mL of USP Methocarbamol RS in *Mobile phase*

Sample solution: 1 mg/mL of Methocarbamol in *Mobile phase*

System suitability

Samples: *System suitability solution* and *Standard solution*

[NOTE—See *Table 1* for relative retention times.]

Suitability requirements

Resolution: NLT 3.5 between methocarbamol and guaifenesin, *System suitability solution*

Tailing factor: NMT 2.0, *Standard solution*

Relative standard deviation: NMT 5.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of each impurity in the portion of Methocarbamol taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (1/F) \times 100$$

r_U = peak response of each impurity from the *Sample solution*

r_S = peak response of methocarbamol from the *Standard solution*

C_S = concentration of USP Methocarbamol RS in the *Standard solution* (mg/mL)

C_U = concentration of Methocarbamol in the *Sample solution* (mg/mL)

F = relative response factor (see *Table 1*)

Acceptance criteria: See *Table 1*.

2 Methocarbamol

Table 1

Name	Relative Retention Time	Relative Response Factor	Acceptance Criteria, NMT (%)
Guaifenesin	0.84	1.2	● post-poned ● (RB 1: May-2015)
Methocarbamol isomer ^a	0.90	1.0	● post-poned ● (RB 1: May-2015)
Methocarbamol	1.0	—	● post-poned ● (RB 1: May-2015)
Methocarbamol dioxolone ^b	1.3	1.0	● post-poned ● (RB 1: May-2015)
Any individual unspecified impurity	—	—	● post-poned ● (RB 1: May-2015)
Total impurities	—	—	2.0

^a 1-Hydroxy-3-(2-methoxyphenoxy)propan-2-yl carbamate.

^b 4-[(2-Methoxyphenoxy)methyl]-1,3-dioxolan-2-one.

▲USP38

SPECIFIC TESTS

- **LOSS ON DRYING** (731)
 Analysis: Dry at 60° for 2 h.
 Acceptance criteria: NMT 0.5%

ADDITIONAL REQUIREMENTS

Change to read:

- **PACKAGING AND STORAGE:** Preserve in tight containers.
 ▲Store at controlled room temperature.▲USP38
- **USP REFERENCE STANDARDS** (11)
 USP Guaifenesin RS
 USP Methocarbamol RS