



Follow the link from the online version of this article to obtain certified continuing medical education credits

Department of Mental Health Sciences, University College London, London W1W 7EJ, UK

Correspondence to: J Carter j.carter@ucl.ac.uk

Cite this as: *BMJ* 2011;343:d5219
doi: 10.1136/bmj.d5219

bmj.com Previous articles in this series

- ▶ Cognitive assessment of older people (*BMJ* 2011;343:d5042)
- ▶ Functional assessment in older people (*BMJ* 2011;343:d4681)
- ▶ Assessing and helping carers of older people (*BMJ* 2011;343:d5202)
- ▶ Fall assessment in older people (*BMJ* 2011;343:d5153)

bmj.com/video

- ▶ Functional assessment in their home

doc2doc

- ▶ Depression—lost in translation. Discuss on BMJ Group's medical community, powered by doc2doc <http://bit.ly/o10012>

Depression in older adults

Joanne Rodda, Zuzana Walker, Janet Carter

Depression is a major contributor to healthcare costs and is projected to be the leading cause of disease burden in middle and higher income countries by the year 2030.^{w1} Depression in later life, traditionally defined as age older than 65, is associated with disability, increased mortality, and poorer outcomes from physical illness. Most clinicians will encounter older patients with depression in their day to day practice, but although treatment is as effective for older patients as for younger adults, the condition is often under-recognised and under-treated. According to WHO data, proportionately more people aged over 65 commit suicide than any other age group, and most have major depression. Older people who attempt suicide are more likely to die than younger people, while in those who survive, prognosis is worse for older adults.¹

With a progressively ageing population worldwide, identification and treatment of depression in older adults becomes increasingly important, especially as older patients may have different presentations and needs than younger ones. We consider recent systematic reviews, meta-analyses, and randomised controlled trials to provide generalists with an understanding of current approaches to the diagnosis and management of patients who develop late life depression.

What is late life depression and who gets it?

Traditionally, the age of 65 has been used to differentiate between “older” and “younger” adults, although there is no set point at which an individual becomes “older” and assessment and care must be based on individual need. Arbitrary definitions of “late life” and differences between studies in terms of diagnostic criteria and populations sampled have produced varying reports of prevalence. Individuals with late life depression represent a hetero-

SOURCES AND SELECTION CRITERIA

We based the review on searches of PubMed, EMBASE, and the Cochrane Database of Systematic Reviews using the search terms “depression”, “elderly”, “aged”, and “old age” published between 2006 and 2011 and limited to English language. We focused on well conducted systematic reviews, meta-analyses, and randomised controlled trials.

geneous group with symptoms that may fall anywhere on a spectrum ranging from sub-threshold mood disorder to major depression. A recent comprehensive meta-analysis using studies with moderate to high methodological quality showed that the point prevalence of major depression in over 75s ranged from 4.6% to 9.3%² whereas rates for sub-threshold depressive symptoms (those failing to reach diagnostic criteria) ranged from 4.5% to 37.4%. A related meta-analysis in people aged over 55 found that sub-threshold depressive symptomatology was two to three times more prevalent than major depression.² Most depressive episodes in late life will be a recurrence rather than a first ever episode^{w2} and the increased female to male ratio is in line with that in younger adults.

Prevalence rates of depression are increased in brain disorders including dementia, Parkinson's disease, and stroke, and also in systemic disease, for example diabetes mellitus and cardiovascular disease (box 1). Prevalence estimates for depression in Alzheimer's disease cluster around 30% but range from 0% to 86%,^{w3} reflecting the difficulty associated with definition and diagnosis of depression in dementia.

How is depression diagnosed in older patients?

Box 2 lists the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) criteria for diagnosis of a major depressive episode. Ideally diagnosis is based on clinical interview, observation of the patient's behaviour, and a collateral history from relatives and care givers. When taking a background history it is important to identify factors that may precipitate and maintain depression. The main risk factors for late life depression are comorbid physical illness, cognitive impairment, functional impairment, lack or loss of close social contacts, and a previous history of depression (box 1), according to the findings of large community based studies.

The risk assessment is important in any psychiatric presentation, and in patients with depression the main area of risk is suicide. Methodologically sound controlled studies have identified some key risk factors for suicide, listed in box 3.

SUMMARY POINTS

- Depression in older adults is associated with an increased risk of death and disability
- Cognitive and functional impairment and anxiety are more common in older than in younger adults with depression
- Older adults with depression are at increased risk of suicide and are more likely than younger adults to complete suicide
- Depression is associated with cognitive impairment and an increased risk of dementia
- A selective serotonin reuptake inhibitor should be the first line pharmacological treatment for depression for most older adults, including those with chronic physical illness
- Psychological and drug treatment is as effective in older as in younger adults
- Subthreshold depressive symptoms that substantially affect older patients' lives are common and management with psychosocial and drug strategies may be effective and prevent further deterioration

Box 1 | Risk factors for depression in elderly people³⁰

Physical factors

Chronic disease, such as diabetes, ischaemic heart disease, heart failure, chronic obstructive pulmonary disease
 Acute myocardial infarction
 Organic brain disease: dementia, stroke, Parkinson's disease, cerebrovascular disease
 Endocrine/metabolic disorders: thyroid disease, hypercalcaemia, B12 and folate deficiency
 Malignancy
 Chronic pain and disability

Psychosocial factors

Social isolation
 Change in financial circumstances
 Being a carer
 Change of role and loss of social status
 Bereavement and loss
 Difficulty in adapting to illness/pain/disability
 Poor defences against anxiety about death
 History of depression
 Being in institutional care

Box 2 | DSM-IV criteria for major depressive episode

Nearly every day for the preceding two weeks the patient has experienced five or more of:
 Depressed mood for most of the day*
 Decreased interest or pleasure in nearly all activities for most of the day*
 Marked loss or gain of weight or markedly increased or decreased appetite
 Excessive sleep or not enough sleep
 Observable psychomotor agitation or retardation
 Tiredness or loss of energy
 Feelings of guilt or worthlessness
 Poor concentration or indecisiveness
 Thoughts of dying or suicide, suicide attempt

*One of these features must be present. Depressed mood for ≥ 2 weeks not meeting these criteria is defined as a minor depressive episode. Diagnostic and statistical manual of mental disorders (DSM-IV), American Psychiatric Association, 1994

Older patients often have symptoms of depression that do not meet the criteria for a major depressive episode (box 2) but are nonetheless clinically important. Identification of the psychological and functional effects of these symptoms determines whether or not treatment is indicated and who may benefit from interventions.

Current guidance for the assessment and management of depression from the UK National Institute for Health and Clinical Excellence (NICE) (<http://guidance.nice.org.uk/CG90/QuickRefGuide/pdf/English>) recommends the use of rating scales to determine severity, although many are weighted towards the presence of somatic symptoms and may therefore overestimate depression in older people, in whom such symptoms are common.

A recent comparison of several assessment scales (patient health questionnaire, Beck depression inventory, hospital anxiety and depression scale) in a primary care population found that treatment and referral rates were identical even though each tool identified differing numbers of patients with moderate to severe depression. However, regardless of the tool used, rates for treatment in older people still remained lower than for younger adults.^{w4} Many rating scales are in common use to assess depression but few are well validated in older people, with the excep-

Box 3 | Risk factors for suicide in older people¹

Older age, male sex
 Social isolation
 Bereavement
 History of attempts
 Evidence of planning
 Chronic painful illness or disability
 Drug or alcohol use
 Sleep disorders

tion of the patient health questionnaire, geriatric depression scale, hospital anxiety and depression scale, and the Cornell scale for depression in dementia (box 4).

Is depression more difficult to diagnose in older adults?

Several studies have shown that older adults are significantly less likely than younger ones to recognise depressive symptoms, which they attribute to normal ageing or physical illness, and that both patients and their doctors tend to view depression as a problem that can be explained away, rather than as an objective illness that warrants treatment.^{w5} These findings suggest that older adults might be less able to identify, and therefore seek appropriate treatment for, common depressive symptoms.

In our clinical experience, late life depression differs qualitatively from depression in early life. Somatisation, hypochondriasis, psychomotor retardation or agitation, and psychosis more commonly form part of the clinical picture, although this tendency has not been uniformly demonstrated.³ Furthermore, late life depression has been associated with cognitive impairment, physical disability, and anxiety, with a large community naturalistic study suggesting that clinically important anxiety coexists in around 50% of patients aged 55-85.⁴

Does depression increase the risk of dementia?

Most studies find that depression in late life is accompanied by measurable cognitive impairment, mediated by memory deficits, diminished executive function, and slowed information processing, which may resolve on remission of symptoms, or may persist even after effective treatment of mood. In the past, the term "depressive pseudodementia" was used to describe reversible dementia in depression, but this oversimplifies the complex spectrum of cognitive impairment.

Two systematic meta-analyses of high quality studies^{5 6} report that late onset depression (after age 65) increases the risk of dementia twofold, but as yet no research has ascertained whether depression is a risk factor for dementia or represents a prodromal condition.⁷

Several mechanisms have been proposed to explain the relation between depression and dementia, including hypercortisolaemia, loss of hippocampal volume, neuro-inflammatory processes, increased Alzheimer-type pathology, reduced cognitive reserve, and vascular disease. None has yet been conclusively demonstrated, but the link is probably multifactorial and the mechanisms not mutually exclusive.

Of these potential mechanisms, vascular changes in the brain have attracted most attention. The key hypothesis is that disruption of prefrontal-striatal circuitry by

Box 4 | Useful scales for depression**Geriatric depression scale (GDS-15)***

Specifically developed for use in geriatric patients; contains fewer somatic items; suitable only for patients with no, mild, or moderate cognitive impairment (>15/30 on mini-mental state examination)

Well validated in older people. Cut-off score in population over 60 of ≥ 5 indicates a case of depression: sensitivity 92%, specificity 54%^{w12 w13}

Cornell scale for depression in dementia (CSDD)*

Suitable for patients with cognitive deficit, not diagnostic for depression but higher scores indicate greater need for further evaluation^{w14}

Patient health questionnaire (PHQ-9)*

Self reported depression assessment tool scoring each of the nine DSM-IV criteria as 0 (not at all) to 3 (nearly every day)

Validated in adults over 60 in primary care in the United States and Netherlands. With cut-off score of ≥ 9 has sensitivity 88%, specificity 80%^{w15 w16}

Beck depression inventory (BDI)

Self reported seven item scale

Not recommended for use in older people owing to focus on somatic symptoms^{w17}

Hospital anxiety and depression scale (HADS)*

Self-rating scale containing two subscales measuring symptoms of depression (HADS-D) and anxiety (HADS-A) during previous week. Scores ≥ 8 for both HADS-A and HADS-D have sensitivity and specificity of 80% and predictive validity of 70%

HADS identifies equal numbers of patients with depression regardless of age^{w18}

Montgomery and Åsberg depression rating scale (MADRS)

Clinician rated 10 item scale, measures severity of depressive symptoms; sensitive to change; mainly used to assess response to treatment but no agreement on cut-off score for remission (between ≤ 4 and ≤ 10), popular in Europe

*Validated in older adults

cerebrovascular pathology produces a syndrome of mood disorder and executive dysfunction. This syndrome is variously described as “vascular depression” or “depression executive dysfunction syndrome”,⁸ reflecting fundamental nosological differences. However, the concept is controversial, and a prospective population based post-mortem study of over 65s found no association between depression and cerebrovascular pathology.⁹ Randomised controlled trials have shown that presence of “vascular depression”/“depression executive dysfunction syndrome” may predict a worse response to antidepressant drug treatment¹⁰ and is associated with increased mortality.¹¹ There is currently no evidence to suggest that treating depression in early or late life reduces the incidence of dementia.

How is late life depression managed?

Given the association between medical morbidity and depression, exclusion of underlying causative or exacer-

bating factors is an important first step in the management of late life depression (box 1). Baseline investigations, for example routine blood tests, may be indicated (box 5).

In subsyndromal and mild depression, psychosocial interventions may be sufficient to cause an improvement. These include increasing social contact and adding structure to the day; for example, assistance in accessing local community events, day centres, or befriending services. Evidence from randomised trials suggests that depressive symptoms in older adults may improve with structured exercise programmes.¹² A RCT of a stepped care approach to the management of subthreshold depressive symptoms found that the intervention (watchful waiting, bibliotherapy based on cognitive behavioural therapy, problem solving therapy, and medication) was associated with a 50% reduction in depression and anxiety disorders at 12 months¹³ compared with treatment as usual and was cost effective.^{w6}

Current NICE guidance recommends that patients with mild or sub-threshold illness who do not respond well to initial supportive interventions are offered psychological therapy or antidepressant medication, while a combination of both interventions is recommended for those with moderate or severe illness.

When should I refer?

NICE guidance recommends that patients are referred to specialist services if they have not responded adequately to management options available in primary care; in severe depression, psychosis, or complex psychosocial situations; and where the degree of risk warrants specialist input. We also emphasise the need to refer older people with comorbid cognitive decline.

Services available in the UK vary geographically and are constantly evolving. A randomised controlled trial of home treatment versus conventional outpatient care for patients aged over 64 living independently and recruited from primary and secondary care services in Austria found significantly reduced depressive symptoms, improved global function, fewer admissions and lower costs of care in the home treatment arm at 3 and 12 months' follow-up.¹⁴ Studies of collaborative care interventions, where care is delivered through integrated mental health and primary care providers, have also repeatedly reported improved outcomes compared with usual care^{w7} although the effect appeared to be associated with prescription of antidepressant medication rather than better communication between primary care providers and mental health services.

Which medication should be prescribed?

Selective serotonin reuptake inhibitors (SSRIs) are well established as first line treatment for depression in older adults. A Cochrane review included 32 randomised controlled trials of antidepressant treatment in people aged 55 or over and reported that SSRIs and tricyclic antidepressants had similar efficacy, but that tricyclics were associated with more side effects and withdrawal from treatment.¹⁵ It was not possible to compare efficacy for other antidepressant groups. Findings from a 2008 meta-analysis of second generation antidepressants in older adults (SSRIs,

Box 5 | Investigations to consider when depression is suspected

Full blood count
Urea and electrolytes
Liver function tests
Thyroid function tests
Vitamin B12
Folate
Fasting glucose
Bone profile
Further tests dictated by clinical presentation

TIPS FOR THE NON-SPECIALIST

- Exclude physical illness as a cause for apparent depressive symptoms
- Bear in mind that factors associated with ageing and the later stages of life, including physical illness, organic brain disease, pain, disability, losses (such as bereavement), and social isolation, create vulnerability to depression
- Be aware that older people with depression may minimise depressive symptoms and may present with somatic problems
- Discuss options for treatment with the patient
- Consider psychosocial interventions first in subsyndromal depressive states and mild depression
- If medication is needed, use an SSRI at a therapeutic dose as first line treatment unless contraindications are present
- Use the same criteria for referral for psychological therapy as in younger adults; older people are just as able to benefit
- Evaluate risk; more people aged over 65 commit suicide than any other age group and most have major depression
- Refer to specialist care if there is substantial risk of self harm, psychosis, need for complex multiprofessional care, inadequate response to treatment, or cognitive impairment
- If treatment is started, evaluate response and need for ongoing treatment regularly

Points to discuss with the patient

- Depression can affect people in different ways; some people may have strong feelings of sadness, but others may be more aware of feeling tired, slowed down, irritable, indecisive, that everything is an effort, or that they worry unnecessarily about small things and experience various physical problems—all these can be symptoms of depression and are not necessarily just part of “getting old”
- There are many different ways to help people get well, for example taking part in social activities, attending clubs and interest groups; physical exercise; talking therapies, and medication
- The beneficial effects of medication may take two to six weeks to be noticeable, but side effects may occur straight away; medication should ideally be continued for at least six months

ADDITIONAL EDUCATIONAL RESOURCES

For patients

Depression (www.ageuk.org.uk/health-wellbeing/conditions-illnesses/depression)—informative web page from Age UK, a charity supporting people in later life

Depression in older adults—(www.rcpsych.ac.uk/mentalhealthinfoforall/problems/depression/depressioninolderadults.aspx) online information leaflet from the UK Royal College of Psychiatrists

CG90 Depression in adults: understanding NICE guidance (<http://guidance.nice.org.uk/CG90/PublicInfo/pdf/English>)—explanation of NICE guidance for those using health services in NHS England and Wales

For healthcare professionals

Depression: the treatment and management of depression in adults (update) (<http://guidance.nice.org.uk/CG90>)—guidance from NICE

Depression (www.cks.nhs.uk/depression/view_whole_topic)—clinical knowledge summary from NHS Evidence

GPNotebook (www.gpnotebook.co.uk)—online medical encyclopaedia

selective serotonin and noradrenaline reuptake inhibitors, bupropion, and mirtazapine) found that treatment in studies lasting 10 weeks or longer was associated with an improved response, supporting the long held belief that response to antidepressants is delayed in older adults.¹⁶ A recent meta-analysis showed an advantage of SSRIs or tricyclic antidepressants over placebo in the treatment of patients with depression in the context of chronic physical illness.¹⁷ Furthermore, evidence from randomised controlled trials has shown that antidepressants are efficacious in depression after stroke¹⁸ and myocardial infarction.¹⁹ Interestingly, a 2007 meta-analysis of 10 randomised

controlled trials of prophylactic antidepressant treatment after stroke reported a significant reduction in the rate of post-stroke depression in treatment groups.²⁰ However, a large randomised controlled trial has recently shown that two commonly used antidepressants, sertraline and mirtazapine, were not appreciably different from placebo in treating depression in patients with Alzheimer’s disease. This effect was sustained at 10 months’ follow-up and side effects were increased in the antidepressant group.²¹

Overall, an SSRI is usually the safest choice in patients with physical illness; the most common drug interactions are mediated via cytochrome p450 enzymes, and citalopram, escitalopram, and sertraline are safest in this regard.

Common side effects of particular concern in the elderly are anticholinergic effects, postural hypotension, and sedation, all of which are more common with tricyclic antidepressants than with SSRIs.²⁰ The risk can be minimised by starting at a low dose and slowly titrating upward. The risk of hyponatraemia induced by antidepressants increases with age and is associated with female sex, low body weight, renal failure, prescription of other drugs associated with hyponatraemia (such as diuretics), and medical comorbidity.^{w8 w9} Older patients prescribed SSRIs are also at increased risk of both upper and lower gastrointestinal bleeding.^{w10} Monitoring of serum sodium levels may be necessary, and the risk of gastrointestinal bleeding can be reduced by prescribing proton pump inhibitors.

NICE guidance recommends that antidepressant treatment is continued for at least six months for a single episode and at least two years if patients are thought to be at risk of relapse. A meta-analysis of eight double-blind placebo controlled trials of maintenance antidepressant therapy between 6 and 36 months in people over 55, published in 2011, found that the optimal duration in older adults is uncertain.²² We suggest that a practical approach is to regularly review depressive symptoms, side effects, comorbidity, and current psychosocial stressors and to involve the patient in the decision making process about ongoing drug treatment.

What if first line drug treatment doesn’t work?

A 2011 systematic review and meta-analysis of inadequate response to treatment in older patients included 13 studies, most of which were open label.²³ The overall response rate for active treatment was 52%, and studies reporting positive results for augmentation of treatment with lithium or antipsychotics, and treatment with venlafaxine, duloxetine, selegiline, or phenelzine, were included. Lithium augmentation was the only treatment for which evidence of efficacy was replicated in more than two studies. We suggest that augmentation of treatment with antipsychotic medication should be used with particular caution in view of the susceptibility of older people to adverse drug reactions, and the paucity of data on safety.

Electroconvulsive therapy is sometimes used after inadequate response to drug treatment, although the usual indication is severe depressive illness in which life threatening refusal of food or fluid, risk of suicide, or psychotic features are present. Electroconvulsive therapy is a safe and effective treatment in the elderly despite an absence of methodologically sound evidence from randomised controlled trials.²⁴

QUESTIONS FOR FUTURE RESEARCH

How can we differentiate between depressive syndromes in older adults, for example those overlapping with anxiety and cognitive impairment?

Does neuroimaging have a role in assessment of depression in older people?

How can we better identify and manage depression in dementia?

Are there ways of preventing depression in older adults at a population level?

What is the optimal period of maintenance treatment for depression in older adults?

Can older adults benefit from psychological therapy?

Results from a 2009 meta-regression analysis suggest that psychological therapy—particularly cognitive behavioural therapy, interpersonal therapy, and problem solving therapy—is equally effective in older and younger adults with depression.²⁵ Combined psychological therapy and pharmacological therapy is more effective than psychological treatment alone for older people with depression.²⁶

What is the outlook for older adults with depression?

A 2005 systematic review of studies comparing outcomes in depression in middle life with those in later life found that rates of remission were similar in both groups, but that late life depression was associated with higher rates of relapse.²⁷ A longitudinal primary care cohort study in the Netherlands reported that the median duration of a major depressive episode in late life was 18 months, with two thirds of patients taking three years to recover.²⁸ In the PRISM-E study, a large study of older patients with major depressive disorder, complete remission was attained in only 29% of patients at six month follow-up.²⁹ Factors associated with prolonged recovery in these studies included severity of depression at baseline, a family history of depression, comorbid anxiety, and general medical comorbidity.

A population based, age stratified, longitudinal study found that adults aged 70-84 years with depression have an increased risk of mortality compared with those who do not have depression, dying on average three years earlier. This risk holds beyond the effects of age, sex, and the presence of dementia, cardiovascular, and other somatic diseases, but did not persist in the oldest old—defined as those aged 85 to 101.^{w11}

Contributors: JR and JC were responsible for the planning, research, writing, and editing of the article. ZW was involved in the planning, writing and editing. JC is the guarantor.

Competing interests: All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Provenance and peer review: Commissioned; externally peer reviewed.

- Manthorpe J, Iliffe S. Suicide in later life: public health and practitioner perspectives. *Int J Geriatr Psychiatry* 2010;25:1230-8.
- Meeks TW, Vahia IV, Lavretsky H, Kulkarni G, Jeste DV. A tune in "a minor" can "b major": a review of epidemiology, illness course, and public health implications of subthreshold depression in older adults. *Affect Disord* 2011;129:126-42.
- Alvarez P, Urretavizcaya M, Benloch L, Vallejo J, Menchon JM. Early- and late-onset depression in the older: no differences found within the melancholic subtype. *Int J Geriatr Psychiatry* 2011;26:615-21.
- Beekman AT, de BE, van Balkom AJ, Deeg DJ, van DR, van TW. Anxiety and depression in later life: Co-occurrence and communality of risk factors. *Am J Psychiatry* 2000;157:89-95.

- Jorm AF. History of depression as a risk factor for dementia: an updated review. *Aust N Z J Psychiatry* 2001;35:776-81.
- Ownby RL, Crocco E, Acevedo A, John V, Loewenstein D. Depression and risk for Alzheimer disease: systematic review, meta-analysis, and meta-regression analysis. *Arch Gen Psychiatry* 2006;63:530-8.
- Byers AL, Yaffe K. Depression and risk of developing dementia. *Nat Rev Neurol* 2011;7:323-31.
- Sneed JR, Culang-Reinlieb ME. The vascular depression hypothesis: an update. *Am J Geriatr Psychiatry* 2011;19:99-103.
- Tsopoulos C, Stewart R, Savva GM, Brayne C, Ince P, Thomas A, et al. Neuropathological correlates of late-life depression in older people. *Br J Psychiatry* 2011;198:109-14.
- Alexopoulos GS, Kiosses DN, Heo M, Murphy CF, Shanmugham B, Gunning-Dixon F. Executive dysfunction and the course of geriatric depression. *Biol Psychiatry* 2005;58:204-10.
- Vilalta-Franch J, Planas-Pujol X, Lopez-Pousa S, Llinas-Regla J, Merino-Aguado J, Garre-Olmo J. Depression subtypes and 5-years risk of mortality in aged 70 years: a population-based cohort study. *Int J Geriatr Psychiatry* 2011, doi:10.1002/gps.2691.
- Blake H, Mo P, Malik S, Thomas S. How effective are physical activity interventions for alleviating depressive symptoms in older people? A systematic review. *Clin Rehabil* 2009;23:873-87.
- Van't Veer-Tazelaar PJ, van Marwijk HW, van Oppen P, van Hout HP, van der Horst HE, Cuijpers P, et al. Stepped-care prevention of anxiety and depression in late life: a randomized controlled trial. *Arch Gen Psychiatry* 2009;66:297-304.
- Klug G, Hermann G, Fuchs-Nieder B, Panzer M, Haider-Stipacek A, Zapotoczky HG, et al. Effectiveness of home treatment for elderly people with depression: randomised controlled trial. *Br J Psychiatry* 2010;197:463-7.
- Mottram P, Wilson K, Strobl J. Antidepressants for depressed elderly. *Cochrane Database Syst Rev* 2006;1:CD003491.
- Nelson JC, Delucchi K, Schneider LS. Efficacy of second generation antidepressants in late-life depression: a meta-analysis of the evidence. *Am J Geriatr Psychiatry* 2008;16:558-67.
- Taylor D, Meader N, Bird V, Pilling S, Creed F, Goldberg D. Pharmacological interventions for people with depression and chronic physical health problems: systematic review and meta-analyses of safety and efficacy. *Br J Psychiatry* 2011;198:179-88.
- Robinson RG, Spalletta G. Poststroke depression: a review. *Can J Psychiatry* 2010;55:341-9.
- Green LA, Dickinson WP, Nease DE, Schellhase KG, Campos-Outcalt D, Schoof BK et al. AAFP guideline for the detection and management of post-myocardial infarction depression. *Ann Fam Med* 2009;7:71-9.
- Chen Y, Patel NC, Guo JJ, Zhan S. Antidepressant prophylaxis for poststroke depression: a meta-analysis. *Int Clin Psychopharmacol* 2007;22:159-66.
- Banerjee S, Hellier J, Dewey M, Romeo R, Ballard C, Baldwin R, et al. Sertraline or mirtazapine for depression in dementia (HTA-SADD): a randomised, multicentre, double-blind, placebo-controlled trial. *Lancet* 2011;378:403-11.
- Kok RM, Heeren TJ, Nolen WA. Continuing treatment of depression in the elderly: a systematic review and meta-analysis of double-blinded randomized controlled trials with antidepressants. *Am J Geriatr Psychiatry* 2011;19:249-55.
- Cooper C, Katona C, Lyketsos K, Blazer D, Brodaty H, Rabins P, et al. A systematic review of treatments for refractory depression in older people. *Am J Psychiatry* 2011.
- Stek ML, Van der Wurff FB, Hoogendijk WL, Beekman AT. Electroconvulsive therapy for the depressed elderly. *Cochrane Database Syst Rev* 2006;2:CD003593.
- Cuijpers P, van SA, Smit F, Andersson G. Is psychotherapy for depression equally effective in younger and older adults? A meta-regression analysis. *Int Psychogeriatr* 2009;21:16-24.
- Cuijpers P, van SA, Warmerdam L, Andersson G. Psychotherapy versus the combination of psychotherapy and pharmacotherapy in the treatment of depression: a meta-analysis. *Depress Anxiety* 2009;26:279-88.
- Mitchell AJ, Subramaniam H. Prognosis of depression in old age compared to middle age: a systematic review of comparative studies. *Am J Psychiatry* 2005;162:1588-601.
- Licht-Strunk E, Van Marwijk HW, Hoekstra T, Twisk JW, De HM, Beekman AT. Outcome of depression in later life in primary care: longitudinal cohort study with three years' follow-up. *BMJ* 2009;338:a3079.
- Azar AR, Chopra MP, Cho LY, Coakley E, Rudolph JL. Remission in major depression: results from a geriatric primary care population. *Int J Geriatr Psychiatry* 2011;26:48-55.
- Colasanti V, Marianetti M, Micacchi F, Amabile GA, Mina C. Tests for the evaluation of depression in the elderly: a systematic review. *Arch Gerontol Geriatr* 2010;50:227-30.

Accepted: 28 July 2011